## THE COVID-19 VACCINE:

## Moderna, Pfizer, Janssen (Johnson & Johnson)

	Moderna	Pfizer	Janssen (J&J)
Safe and Effective?	•	<b>*</b>	
Side Effects	83.9% - 90.5% of participants had short-term pain at the injection site, and 71.9% - 81.9% of participants experienced side effects	87% of participants had short-term pain at the injection site and 77% of participants experienced side effects	<b>48.6%</b> of participants hadshort-term pain at the injection site, and 33.2% -38.9% experienced side effects (e.g. fever, fatigue, headache, chills)
Large Clinical Trial Size	<b>30,000</b> participant randomized, doubleblind, placebo-controlled trial	<b>43,000</b> participant randomized, doubleblind, placebo-controlled trial	<b>44,000</b> participant randomized, double-blind, placebo-controlled trial
Storage and Handling	Stored in freezer, but stable in refrigerator for 30 days	Stored in ultra-cold freezers	Stored in temperatures 36 to 46 degrees Fahrenheit
Dosing	(0.5 ml) - two doses, 28 days apart Moderna or Pfizer Booster needed	(0.3 ml/.15 pediatric) - two doses, 21 days apart Moderna or Pfizer Booster needed	(0.5 ml) single dose Moderna or Pfizer Booster needed*
Age Range	<b>18 years</b> and older	<b>5 years</b> and older 5-17 year olds must be accompanied by a parent or guardian	<b>18 years</b> and older



All Vaccines are free and available! 2507 Wahnish Way, Tallahassee, FL 32307 8 a.m. to 4:30 p.m. Monday through Saturday

<sup>\*</sup>According to the CDC, boosters of an mRNA vaccine are preferred over the Johnson & Johnson vaccine (Dec. 2021).